



TOSOH



Analys & Mätteknik

031 336 90 00 • www.scantecnordic.se

ANALYSIS



TSKgel® SuperSW mAb HTP/HR TSKgel® UltraSW AGGREGATE

Aqueous size exclusion chromatography (SEC) is the method of choice for the analysis of protein fragments, monomers, and aggregates under non-denaturing conditions. Based on the flow of the sample through a porous stationary phase SEC separates molecules according to their size, or more precisely, their hydrodynamic volume. In aqueous elution systems SEC is also referred to as gel filtration chromatography (GFC). TSKgel G3000SW_{XL} columns have been the industry's standard for quality control of monoclonals by SEC for decades. Based on the proven proprietary surface technology of the renowned TSKgel SW series, a new series of silica-based SEC columns was engineered to provide shorter analysis time or higher resolution for antibody analysis.

HIGHLIGHTS


- Optimized for antibody analysis
- Small particle size for UHPLC use
- Highest resolution with SuperSW mAb HR
- Fast separation with SuperSW mAb HTP
- Higher molecular weight range with UltraSW Aggregate

FEATURES

The new series of dedicated SEC columns for mAb analysis delivers significant advancements in resolution. It comprises of three different columns and their guards. Table 1 summarizes the characteristics of the mAb SEC columns and Figure 1 shows the calibration curves and the molecular weight range of the three columns.

The calibration curve of TSKgel SuperSW mAb HR is most similar to the one of TSKgel G3000SW_{XL}, the current industrial standard for antibody analysis. The new columns are highly reproducible. The high Batch-to-Batch stability is proved in Figure 2 for TSKgel SuperSW mAb HR. The lifetime of the columns can be further improved when using the corresponding guard columns.

CALIBRATION CURVE



➤ Figure 1

Columns: TSKgel SuperSW mAb HTP 4.6 mm ID × 15 cm, TSKgel SuperSW mAb HR, TSKgel UltraSW Aggregate, TSKgel G3000SW_{XL} (all 7.8 mm ID × 30 cm)

Mobile phase: 0.2 mol/L phosphate buffer, pH 6.7 0.05% Na₂S₂O₃; Flow rate: 1.0 mL/min, 0.35 mL/min (SuperSW mAb HTP); Temp.: 25°C; Detection: UV 280 nm; Inj. vol.: 10 µL, 5 µL (SuperSW mAb HTP)

1. Thyroglobulin (MW 640,000), 2. γ-Globulin (MW 155,000),
3. Ovalbumin (MW 47,000), 4. Ribonuclease A (MW 13,700)
5. p-Aminobenzoic acid (MW 137)

CHARACTERISTICS OF TSKgel mAb COLUMNS

| Column | TSKgel SuperSW mAb HR | TSKgel SuperSW mAb HTP | TSKgel UltraSW Aggregate |
|--------------------------------------|-----------------------|------------------------|--------------------------|
| Dimension | 7.8 mm ID x 30 cm | 4.6 mm ID x 15 cm | 7.8 mm ID x 30 cm |
| Theoretical plates | ≥ 30.000 | ≥ 15.000 | ≥ 35.000 |
| Base material | Silica gel | | Silica gel |
| Particle size | 4 µm | | 3 µm |
| Separation range (globular proteins) | 10,000 - 500,000 Da | | 10,000 - 2,000,000 Da |

➤ Table 1

Each of the new SEC columns is tailored to a specific separation problem. TSKgel SuperSW mAb HTP - "HTP" standing for high throughput - was developed to enable an easy transfer of HPLC methods based on TSKgel SW_{XL} to fast UHPLC analysis. Small particle size silica beads are packed in UHPLC column hardware with 4.6 mm inner diameter. This enables to double the throughput without compromising resolution (Figure 3).

TSKgel SuperSW mAb HR - "HR" indicating high resolution - delivers superior resolution over the whole range from fragments to aggregates when analysing monoclonals (Figure 4). It has the same column dimensions - 7.8 mm by 30 cm length - as the established TSKgel G3000SW_{XL}. TSKgel Ultra SW Aggregate (7.8 mm ID x 30 cm) features a smaller particle size and a higher exclusion limit. It offers a wider separation window in the aggregate region (Figure 4 C).

BATCH TO BATCH REPRODUCIBILITY




Figure 2
 Column: TSKgel SuperSW mAb HR (7.8 mm ID x 30 cm)
 Eluent: 0.2 mol/L phosphate buffer (pH 6.7)+ 0.05% NaN₃
 Flow rate: 0.8 mL/min; Detection: UV @ 280 nm; Inj. volume: 10 µL;
 Sample: 1. Thyrobulobulin, 2.γ-Globulin, 3. Ovalbumin, 4. Ribonuclease A, 5. p-Aminobenzoic acid

Ordering information

| Part-No | Description | Matrix | Housing | Dimensions |
|---------|---------------------------------------|--------|-----------------|-----------------------|
| 0022854 | TSKgel SuperSW mAb HR, 4 µm, 25 nm | Silica | Stainless steel | 7.8 mm ID x 30.0 cm L |
| 0022855 | TSKgel SuperSW mAb HTP, 4 µm, 25 nm | Silica | Stainless steel | 4.6 mm ID x 15.0 cm L |
| 0022856 | TSKgel UltraSW Aggregate, 3 µm, 30 nm | Silica | Stainless steel | 7.8 mm ID x 30.0 cm L |
| 0022857 | TSKgel Guardcolumn SuperSW mAb, 4 µm | Silica | Stainless steel | 6.0 mm ID x 4.0 cm L |
| 0022858 | TSKgel Guardcolumn SuperSW mAb, 4 µm | Silica | Stainless steel | 3.0 mm ID x 2.0 cm L |
| 0022859 | TSKgel Guardcolumn UltraSW, 3 µm | Silica | Stainless steel | 6.0 mm ID x 4.0 cm L |

FAST ANALYSIS OF mAb AGGREGATION




Figure 3
 Column: TSKgel SuperSW mAb HTP (4.6 mm ID x 15 cm)
 Elution: 0.2 mol/L phosphate buffer (pH 6.7) + 0.05% NaN₃
 Flow rate: 0.50 mL/min, 0.35 mL/min; Detection: UV @ 280 nm
 Temp.: 25°C; Sample: monoclonal antibody (mouse-human chimeric IgG, Erbitux), 5 µL

COMPARISON OF AGGREGATE ANALYSIS




Figure 4
 Columns: A. TSKgel G3000SW_{XL}, B. TSKgel SuperSW mAb HR, C. TSKgel UltraSW Aggregate; Dimension: 7.8 mm ID x 30 cm;
 Eluent: 0.2 mol/L phosphate buffer (pH 6.7) + 0.05% NaN₃
 Flow rate: 0.8 mL/min; Detection: UV @ 280 nm; Temp.: 25°C;
 Sample: monoclonal antibody, (mouse-human chimeric IgG, Erbitux), 10 µL